

Balloons and Hoops and their Universal Finite-Type Invariant,

BF Theory, and an Ultimate Alexander Invariant

Dror Bar-Natan in Hamburg, August 2012

$\omega \in \beta := \text{http://www.math.toronto.edu/~drorbn/Talks/Hamburg-1208}$




Scheme. • Balloons and hoops in \mathbb{R}^4 , algebraic structure and relations with 3D.



• An ansatz for a “homomorphic” invariant: computable, related to finite-type and to BF.

• Reduction to an “ultimate Alexander invariant”.

$\mathcal{K}^{bh}(m, n)$.



Examples.




I mean business!

```

S_0 = Rm[3, a] Rp[b, 2] Sp[1, 4];
S = S_0 // dm[2, 1, 1] // dm[4, b, b] //
      dm[1, a, a] // dm[3, a, a];
S[[5]] /. {w_LW -> (Deg[w] + 1) * w,
          w_CW -> Deg[w] * w}

mu[CWS[-[a], -2 [ab], -3 [aab] - 3 [abb],
-4 [aaab] + 42 [aabb] - 60 [ababb] - 4 [abbbb],
-5 [aaaa] + 110 [aaabb] - 180 [aabbab] -
110 [aabbba] - 180 [ababbb] - 5 [abbbb]],
h[b] LS[2 (a, 0, -24 (aab),
-60 (aaab) + 60 (aabb), -120 (aaaab) +
900 (aaabb) - 360 (aabbab) - 120 (aabbba)] -
h[a] LS[-2 (a + 2 (b), 9 (ab), 26 (aab) -
26 (abb), 60 (aaab) - 255 (aabb) + 60 (abbbb),
119 (aaaab) - 1504 (aaabb) + 118 (aabbab) -
1504 (aabbba) - 1386 (ababbb) - 119 (abbbb)]]
    
```

Tangles (u/v/w).



• δ injects u-Knots into \mathcal{K}^{bh} (likely u-tangles too).
 • δ maps v/w-tangles map to \mathcal{K}^{bh} ; the kernel contains Reidemeister moves and the “overcrossings commute” relation, and conjecturally, that’s all. Allowing punctures and cuts, δ is onto.

Operations
Punctures & Cuts



Meta-Group-Action.

If X is a space, $\pi_1(X)$ is a group, $\pi_2(X)$ is an Abelian group, and π_1 acts on π_2 .

“MGA”




(“//” is newspeak for “apply an operator” and for “composition left to right”)


Properties.

- Associativities: $m_a^{ab} // m_a^{ac} = m_b^{bc} // m_a^{ab}$, for $m = tm, hm$.
- Action axiom t : $tm_w^{uv} // hta^{xw} = hta^{xu} // hta^{xv} // tm_w^{uv}$.
- Action axiom h : $hm_z^{xy} // hta^{zu} = hta^{xu} // hta^{yu} // hm_z^{xy}$.
- SD Product: $dm_c^{ab} := hta^{ba} // tm_c^{ab} // hm_c^{ab}$ is associative.

Meta-associativity.



Tangle concatenations $\rightarrow \pi_1 \times \pi_2$.



Thus we seek homomorphic invariants of \mathcal{K}^{bh} !

Invariant #0. With Π_1 denoting “honest π_1 ”, map $\gamma \in \mathcal{K}^{bh}(m, n)$ to the triple $(\Pi_1(\gamma^c), (u_i), (x_j))$, where the meridian of the balls u_i normally generate Π_1 , and the “longitudes” x_j are some elements of Π_1 .
 * acts like *, tm acts by “merging” two meridians/generators, hm acts by multiplying two longitudes, and hta^{xu} acts by “conjugating a meridian by a longitude”:

$(\Pi, (u, \dots), (x, \dots)) \mapsto (\Pi * \langle \bar{u} \rangle / (u = x \bar{u} x^{-1}), (\bar{u}, \dots), (x, \dots))$

Failure #0. Can we write the x ’s as free words in the u ’s?
 If $x = uv$, compute $x // hta^{xu}$:

$$x = uv \rightarrow \bar{u}v = u^x v = u^{\bar{u}v} v = u^{u^x v} v = u^{u^x v} v = \dots$$

The Meta-Group-Action M. Let T be a set of “tail labels” (“balloon colours”), and H a set of “head labels” (“hoop colours”).

Let $FL = FL(T)$ and $FA = FA(T)$ be the (completed graded) free Lie and free associative algebras on generators T and let $CW = CW(T)$ be the (completed graded) vector space of cyclic words on T , so there’s $\text{tr} : FA \rightarrow CW$.
 Let $M(T, H) := \{(\bar{\lambda} = (x : \lambda_x)_{x \in H}; \omega) : \lambda_x \in FL, \omega \in CW\}$

$$= \left\{ \left(x : \begin{array}{c} u \\ \diagdown \\ v \end{array}, y : \begin{array}{c} v \\ \diagup \\ u \end{array} \mid -\frac{22}{7} \begin{array}{c} u \\ \diagup \\ v \end{array} \begin{array}{c} u \\ \diagdown \\ v \end{array}; \begin{array}{c} u \\ \diagdown \\ v \end{array} \begin{array}{c} v \\ \diagup \\ u \end{array} \right) \dots \right\}$$


Operations. Set $(\bar{\lambda}_1; \omega_1) * (\bar{\lambda}_2; \omega_2) := (\bar{\lambda}_1 \cup \bar{\lambda}_2; \omega_1 + \omega_2)$ and with $\mu = (\bar{\lambda}; \omega)$ define

$$tm_w^{uv} : \mu \mapsto \mu // (u, v \mapsto w),$$

$$hm_z^{xy} : \mu \mapsto \left(\left(\dots, x : \lambda_x, y : \lambda_y, \dots, z : \text{bch}(\lambda_x, \lambda_y) \right); \omega \right)$$

$$hta^{xu} : \mu \mapsto \underbrace{\mu // \langle \bar{u} \rangle}_{\mu // CC_u^{\lambda_x}} (u \mapsto e^{\text{ad } \lambda_x}(\bar{u})) // (\bar{u} \mapsto u) + \underbrace{(0; J_u(\lambda_x))}_{\text{the “J-spice”}}$$

A CC_u^λ example.




Balloons and Hoops and their Universal Finite-Type Invariant, 2

The Meta-Cocycle J . Set $J_u(\lambda) := J(1)$ where

$$J(0) = 0, \quad \lambda_s = \lambda // CC_u^{s\lambda},$$

$$\frac{dJ(s)}{ds} = (J(s) // \text{der}(u \mapsto [\lambda_s, u])) + \text{div}_u \lambda_s,$$

and where $\text{div}_u \lambda := \text{tr}(u\sigma_u(\lambda))$, $\sigma_u(v) := \delta_{uv}$, $\sigma_u([\lambda_1, \lambda_2]) := \iota(\lambda_1)\sigma_u(\lambda_2) - \iota(\lambda_2)\sigma_u(\lambda_1)$ and ι is the inclusion $FL \hookrightarrow FA$:



Claim. $CC_u^{\text{bch}(\lambda_1, \lambda_2)} = CC_u^{\lambda_1} // CC_u^{\lambda_2} // CC_u^{\lambda_1}$ and


$$J_u(\text{bch}(\lambda_1, \lambda_2)) = J_u(\lambda_1) // CC_u^{\lambda_2} // CC_u^{\lambda_1} + J_u(\lambda_2 // CC_u^{\lambda_1}),$$

and hence tm , hm , and hta form a meta-group-action.

Why ODEs? **Q.** Find f s.t. $f(x+y) = f(x)f(y)$.

A. $\frac{df(s)}{ds} = \frac{d}{de} f(s + \epsilon) = \frac{d}{de} f(s)f(\epsilon) = f(s)C$.

Now solve this ODE using Picard's theorem or power series.



The β quotient, 2. Let $R = \mathbb{Q}[\{c_u\}_{u \in T}]$ and $L_\beta := R \otimes T$ with central R and with $[u, v] = c_u v - c_v u$ for $u, v \in T$. Then $FL \rightarrow L_\beta$ and $CW \rightarrow R$. Under this,

$$\mu \rightarrow (\bar{\lambda}; \omega) \quad \text{with } \bar{\lambda} = \sum_{x \in H, u \in T} \lambda_{ux} u x, \quad \lambda_{ux}, \omega \in R,$$

$$\text{bch}(u, v) \rightarrow \frac{c_u + c_v}{e^{c_u + c_v} - 1} \left(\frac{e^{c_u} - 1}{c_u} u + e^{c_u} \frac{e^{c_v} - 1}{c_v} v \right),$$

if $\lambda = \sum \lambda_v v$ then with $c_\lambda := \sum \lambda_v c_v$,

$$u // CC_u^\lambda = \left(1 + c_u \lambda_u \frac{e^{c_\lambda} - 1}{c_\lambda} \right)^{-1} \left(e^{c_\lambda} u - c_u \frac{e^{c_\lambda} - 1}{c_\lambda} \sum_{v \neq u} \lambda_v v \right),$$

$\text{div}_u \lambda = c_u \lambda_u$, and the ODE for J integrates to

$$J_u(\lambda) = \log \left(1 + \frac{e^{c_\lambda} - 1}{c_\lambda} c_u \lambda_u \right),$$

so ζ is formula-computable to all orders! **Can we simplify?**

Repackaging. Given $((x : \lambda_{ux}); \omega)$, set $c_x := \sum_v c_v \lambda_{vx}$, replace $\lambda_{ux} \rightarrow \alpha_{ux} := c_u \lambda_{ux} \frac{e^{c_x} - 1}{c_x}$ and $\omega \rightarrow \log \omega$, use $t_u = e^{c_u}$, and write α_{ux} as a matrix. Get " **β calculus**".

The Invariant ζ . Set $\zeta(\rho^\pm) = (\pm u_x; 0)$. This at least defines an invariant of u/v/w-tangles, and if the topologists will deliver a "Reidemeister" theorem, it is well defined on \mathcal{K}^{bh} .

$$\zeta: \begin{array}{c} \text{u} \\ \text{v} \\ \text{w} \end{array} \xrightarrow{\text{bch}} (x : + |^u ; 0) \quad \begin{array}{c} \text{u} \\ \text{v} \\ \text{w} \end{array} \xrightarrow{\text{bch}} (x : - |^u ; 0)$$

Theorem. ζ is (the log of) a universal finite type invariant (a homomorphic expansion) of w-tangles.

Tensorial Interpretation. Let \mathfrak{g} be a finite dimensional Lie algebra (any!). Then there's $\tau : FL(T) \rightarrow \text{Fun}(\oplus_T \mathfrak{g} \rightarrow \mathfrak{g})$ and $\tau : CW(T) \rightarrow \text{Fun}(\oplus_T \mathfrak{g})$. Together, $\tau : M(T, H) \rightarrow \text{Fun}(\oplus_T \mathfrak{g} \rightarrow \oplus_H \mathfrak{g})$, and hence

$$e^\tau : M(T, H) \rightarrow \text{Fun}(\oplus_T \mathfrak{g} \rightarrow \mathcal{U}^{\otimes H}(\mathfrak{g})).$$

ζ and BF Theory. Let A denote a \mathfrak{g} -connection on S^4 with curvature F_A , and B a \mathfrak{g}^* -valued 2-form on S^4 . For a hoop γ_x , let $\text{hol}_{\gamma_x}(A) \in \mathcal{U}(\mathfrak{g})$ be the holonomy of A along γ_x . For a ball γ_u , let $\mathcal{O}_{\gamma_u}(B) \in \mathfrak{g}^*$ be the integral of B (transported via A to ∞) on γ_u .



Loose Conjecture. For $\gamma \in \mathcal{K}(T, H)$,

$$\int \mathcal{D}A \mathcal{D}B e^{\int B \wedge F_A} \prod_u e^{\mathcal{O}_{\gamma_u}(B)} \bigotimes_x \text{hol}_{\gamma_x}(A) = e^\tau(\zeta(\gamma)).$$

That is, ζ is a complete evaluation of the BF TQFT.

Issues. How exactly is B transported via A to ∞ ? How does the ribbon condition arise? Or if it doesn't, could it be that ζ can be generalized??

The β quotient, 1. • Arises when \mathfrak{g} is the 2D non-Abelian Lie algebra.

• Arises when reducing by relations satisfied by the weight system of the Alexander polynomial.

β Calculus. Let $\beta(H, T)$ be

$$\left\{ \begin{array}{c|ccc} \omega & x & y & \cdots \\ \hline u & \alpha_{ux} & \alpha_{uy} & \cdot \\ v & \alpha_{vx} & \alpha_{vy} & \cdot \\ \vdots & \cdot & \cdot & \cdot \end{array} \middle| \begin{array}{l} \omega \text{ and the } \alpha_{ux}'\text{s are} \\ \text{rational functions in} \\ \text{variables } t_u, \text{ one for} \\ \text{each } u \in T. \end{array} \right\},$$



In preparation, Selmani & B-N.

$$tm_w^{uv} : \begin{array}{c|ccc} \omega & \cdots \\ \hline u & \alpha & \beta \\ v & \beta & \gamma \\ \vdots & \vdots & \vdots \end{array} \mapsto \begin{array}{c|ccc} \omega & \cdots \\ \hline w & \alpha + \beta & \gamma \end{array}, \quad \begin{array}{c|cc} \omega_1 & H_1 \\ \hline T_1 & \alpha_1 \end{array} \cup \begin{array}{c|cc} \omega_2 & H_2 \\ \hline T_2 & \alpha_2 \end{array} = \begin{array}{c|ccc} \omega_1 \omega_2 & H_1 & H_2 \\ \hline T_1 & \alpha_1 & 0 \\ T_2 & 0 & \alpha_2 \end{array},$$

$$hm_z^{xy} : \begin{array}{c|ccc} \omega & x & y & \cdots \\ \hline \vdots & \alpha & \beta & \gamma \end{array} \mapsto \begin{array}{c|ccc} \omega & & & \cdots \\ \hline \vdots & \alpha + \beta + \langle \alpha \rangle \beta & \gamma & \end{array},$$

$$hta^{xu} : \begin{array}{c|ccc} \omega & x & \cdots \\ \hline u & \alpha & \beta \\ \vdots & \gamma & \delta \end{array} \mapsto \begin{array}{c|ccc} \omega \epsilon & & & \cdots \\ \hline u & \alpha(1 + \langle \gamma \rangle / \epsilon) & \beta(1 + \langle \gamma \rangle / \epsilon) \\ \vdots & \gamma / \epsilon & \delta - \gamma \beta / \epsilon \end{array},$$

where $\epsilon := 1 + \alpha$, $\langle \alpha \rangle := \sum_v \alpha_v$, and $\langle \gamma \rangle := \sum_{v \neq u} \gamma_v$, and let

$$R_{ux}^+ := \frac{1}{u} \left| \begin{array}{c} x \\ t_u - 1 \end{array} \right. \quad R_{ux}^- := \frac{1}{u} \left| \begin{array}{c} x \\ t_u^{-1} - 1 \end{array} \right.$$

On long knots, ω is the Alexander polynomial!

Why bother? (1) An ultimate Alexander invariant: Manifestly polynomial (time and size) extension of the (multi-variable) Alexander polynomial to tangles. Every step of the computation is the computation of the invariant of some topological thing (no fishy Gaussian elimination!). *If there should be an Alexander invariant to have an algebraic categorification, it is this one!* See also $\omega \epsilon \beta / \text{regina}$, $\omega \epsilon \beta / \text{gwu}$.

Why bother? (2) Related to A-T, K-V, and E-K, should have vast generalization beyond w-knots and the Alexander polynomial. See also $\omega \epsilon \beta / \text{wko}$, $\omega \epsilon \beta / \text{caen}$, $\omega \epsilon \beta / \text{swiss}$.



"God created the knots, all else in topology is the work of mortals."
Leopold Kronecker (modified)



Balloons and Hoops and their Universal Finite-Type Invariant, 3

Abstract. Balloons are two-dimensional spheres. Hoops are one dimensional loops. Knotted Balloons and Hoops (KBH) in 4-space behave much like the first and second fundamental groups of a topological space - hoops can be composed like in π_1 , balloons like in π_2 , and hoops “act” on balloons as π_1 acts on π_2 . We will observe that ordinary knots and tangles in 3-space map into KBH in 4-space and become amalgams of both balloons and hoops.

We give an ansatz for a tree and wheel (that is, free-Lie and cyclic word) -valued invariant Z of KBHs in terms of the said compositions and action and we explain its relationship with finite type invariants. We speculate that Z is a complete evaluation of the BF topological quantum field theory in 4D, though we are not sure what that means. We show that a certain “reduction and repackaging” of Z is an “ultimate Alexander invariant” that contains the Alexander polynomial (multivariable, if you wish), has extremely good composition properties, is evaluated in a topologically meaningful way, and is least-wasteful in a computational sense. If you believe in categorification, here’s a wonderful playground.

